



Carrie Adams Lab
Restoration and Plant Ecology
Recent work in the region

- Plant focused research on ecosystem functions
- Genetic diversity of *Juncus* roemerianus
- Salinity and flooding stress thresholds of salt marsh plants
- Planting guidance for Living Shorelines
- Patterns of stress in *Juncus* marshes





Project motivations

- Observed death of coastal forest trees and reduced regeneration of seedlings
- Substantial marsh loss in the Northeastern US and Louisiana
- Current marsh conditions along
 Florida's Gulf coast are understudied

Project Objectives



Quantify

Juncus roemerianus morphological and physiological conditions as possible indicators of stress leading to die-off in the Big Bend region



Assess


Juncus conditions across spatial locations within marsh systems



Examine


Patterns in *Juncus* response across environmental factors (soil, water, and herbivory) to determine within- and between- marsh variation

















Salt marsh parameters

Juncus conditions

- Productivity (biomass)
- Stem height and density
- Tissue enzyme (proline) and ion accumulation
- Plant water conductance (stomate density)


Environmental factors

- Soil and groundwater salinity
- Soil nutrient levels (C and N)
- Herbivore presence (snail density)

Objective 1: Quantifying *Juncus* conditions


Evidence of stress present in 8 of the 15 measured response variables

- Physiological responses point to Juncus experiencing stress
- Productivity did **not** show strong indication of *Juncus* stress










Objective 2: Assess *Juncus* conditions across spatial locations















Objective 3: Examine *Juncus* response to environmental variables to determine withinand between- marsh variation

- Plant stress varied considerably between marsh systems
 - Proximity to Suwannee River showed fewest signs of stress
- Soils varied in texture, organic matter, nutrients, and bulk density
- Groundwater salinity seasonal trends across creek systems
 - Highest salinities in dry season
 - Lowest salinities in wet season

Project conclusions

- Physiological adaptations buffer *Juncus* from stressful environmental conditions
- Future marsh loss (conversion to open water) to occur at creek margins
- Florida's Big Bend salt marshes are at various stages of threat to marsh die-off
- Fresh water river discharge and rain events likely reduce salinity stress



Thank you!

Special thanks:

Cher Nicholson Sarah Tevlin Natalia Medina Alex Matys

Dr. Carrie Adams

Dr. Bill Pine







