

ISSN 1930-1448

Oyster Integrated Mapping and Monitoring Program Report for the State of Florida

KARA R. RADABAUGH, STEPHEN P. GEIGER, RYAN P. MOYER, EDITORS



Florida Fish and Wildlife Conservation Commission
Fish and Wildlife Research Institute
Technical Report No. 22 • 2019
MyFWC.com

Oyster Integrated Mapping and Monitoring Program Report for the State of Florida

KARA R. RADABAUGH, STEPHEN P. GEIGER, RYAN P. MOYER, EDITORS



Florida Fish and Wildlife Conservation Commission
Fish and Wildlife Research Institute
100 Eighth Avenue Southeast
St. Petersburg, Florida 33701
MyFWC.com

Technical Report 22 • 2019



Ron DeSantis
Governor of Florida



Eric Sutton
Executive Director
Florida Fish and Wildlife Conservation Commission

The [Fish and Wildlife Research Institute](#) is a division of the Florida Fish and Wildlife Conservation Commission, which “[manages] fish and wildlife resources for their long-term well-being and the benefit of people.” The Institute conducts applied research pertinent to managing fishery resources and species of special concern in Florida. Programs focus on obtaining the data and information that managers of fish, wildlife, and ecosystems need to sustain Florida’s natural resources. Topics include managing recreationally and commercially important fish and wildlife species; preserving, managing, and restoring terrestrial, freshwater, and marine habitats; collecting information related to population status, habitat requirements, life history, and recovery needs of upland and aquatic species; synthesizing ecological, habitat, and socioeconomic information; and developing educational and outreach programs for classroom educators, civic organizations, and the public.

The Institute publishes three series: *Memoirs of the Hourglass Cruises*, *Florida Marine Research Publications*, and *FWRI Technical Reports*. *FWRI Technical Reports* contain information relevant to immediate needs in resource management.

Gil McRae, FWRI Director
Bland Crowder, Editor and Production

This publication is available online and may be downloaded at <http://myfwc.com/research/publications/scientific/technical-reports/>.

We suggest that this document be cited as follows:

Radabaugh KR, Moyer RP, Geiger SP, editors. 2019. Oyster integrated mapping and monitoring program report for the state of Florida. St. Petersburg, FL: Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission. FWRI Technical Report 22.

Cover and text papers used in this publication meet the minimum requirements of the American National Standard for Permanence of Paper for Printed Library Materials Z39.48–1992(R2009).

Cover photograph: An eastern oyster (*Crassostrea virginica*) reef in St. Petersburg, Florida. Photograph by Kara Radabaugh.

Contents

Acknowledgments	<u>iv</u>
Summary of the Oyster Integrated Mapping and Monitoring Program	<u>v</u>
OIMMP Report Authors and Contributors.....	<u>vi</u>
Executive Summary	<u>vii</u>
Chapter 1: Introduction	<u>1</u>
Introduction to oysters in Florida	<u>1</u>
Eastern oyster life history and ecology	<u>2</u>
Oyster reefs and shell budgets	<u>4</u>
Oyster harvesting in Florida	<u>5</u>
Threats to oyster reefs	<u>7</u>
Mapping oyster reefs	<u>12</u>
Recent oyster mapping data in Florida	<u>17</u>
Oyster reef monitoring.....	<u>18</u>
Region-specific chapters	<u>23</u>
Chapter 2: Northwest Florida	<u>35</u>
Chapter 3: Apalachicola Bay	<u>48</u>
Chapter 4: Big Bend and Springs Coast	<u>68</u>
Chapter 5: Tampa and Sarasota Bays	<u>83</u>
Chapter 6: Southwest Florida	<u>96</u>
Chapter 7: Southeast Florida	<u>116</u>
Chapter 8: Central Eastern Florida.....	<u>124</u>
Chapter 9: Northeast Florida	<u>143</u>
Chapter 10: Conclusions and Recommendations	<u>160</u>
Appendix A: Historical Yields of the Florida Oyster Fishery	<u>164</u>
Appendix B: Abbreviations	<u>171</u>
Appendix C: Species List	<u>174</u>

Acknowledgments

This report was funded by grants from the U.S. Fish and Wildlife Service in support of Florida's State Wildlife Grants Program (Award numbers F16AF00528 and F19AF00401), administered by the Florida Fish and Wildlife Conservation Commission (FWC) to support the study of species of greatest conservation need and high-priority coastal habitats in accordance with the Florida State Wildlife Action Plan. The Oyster Integrated Mapping and Monitoring Program (OIMMP) editors wish to thank Andrea Alden, Caroline Gorga, Melanie Parker, Kathleen O'Keefe, and Amber Whittle for their support and guidance. We also wish to thank the dozens of scientists and managers across Florida who contributed directly to the writing of this report and the many others who provided comments and contributions during the OIMMP workshops, hosted by Nikki Dix and Kaitlyn Dietz at the Guana Tolomato Matanzas National Estuarine Research Reserve. Christi Santi and Kara Radabaugh created region-specific maps and graphs for this report. Amber Whittle and William Arnold completed technical review of the document. Bland Crowder completed scientific review, copy review, and layout. Research, mapping, and monitoring support were provided by the staff of the Coastal Wetlands Research Program at FWC's Fish and Wildlife Research Institute, including Amanda Chappel, Emma Dontis, Scott Adams, Christine Russo, Sam Roussopoulos, Joshua Breithaupt, Sierra Greene, and Emma Wennick.

This collaborative report has benefited from the contributions of many authors from numerous governmental and independent agencies. The views, statements, findings, conclusions, and recommendations expressed herein are those of the authors and do not necessarily reflect the views of the State of Florida, the Florida Water Management Districts, the National Oceanic and Atmospheric Administration, or any of their subagencies.



An assortment of oysters on red mangrove (*Rhizophora mangle*) prop roots in the Ten Thousand Islands, Florida. Photo credit: Kara Radabaugh.

Summary of the Oyster Integrated Mapping and Monitoring Program

The Oyster Integrated Mapping and Monitoring Program (OIMMP) began as a joint effort between the Coastal Wetlands and Molluscan Fisheries research programs at the Florida Fish and Wildlife Conservation Commission's Fish and Wildlife Research Institute in St. Petersburg, Florida. OIMMP is based on the framework established by the Seagrass Integrated Mapping and Monitoring (SIMM) program (<http://myfwc.com/research/habitat/seagrasses/projects/active/simm/>) and the Coastal Habitat Integrated Mapping and Monitoring Program (CHIMMP) (<http://myfwc.com/research/habitat/coastal-wetlands/projects/chimmp/>), which rely upon a network of ecosystem experts to assemble regional summaries of mapping and monitoring data. The main objective of OIMMP was to build and maintain a collaborative network of stakeholders with interest in mapping and monitoring Florida's oyster habitats in order to identify the status of and management priorities for oysters and their habitats.

OIMMP workshops were held at the Guana Tolomato Matanzas National Estuarine Research Reserve in 2017 and 2018 and the Fish and Wildlife Research Institute in 2019 to bring together oyster researchers and managers from across the state. During these workshops, attendees gave presentations on oyster mapping and monitoring activities and made recommendations for future mapping, monitoring, and management of oyster resources. See <http://ocean.floridamarine.org/OIMMP> for detailed proceedings and outcomes of the OIMMP workshops.

Attendees of the 2017 workshop developed the regional boundaries for the chapters in this report, and many attendees also volunteered to contribute their expertise as coauthors. Additional regional coauthors were added based on need and personal recommendations (see below for a list of all regional contributors). Due to the collaborative nature of this report, the style, content, and level of detail varies among chapters based upon regional data availability, range of participating organizations, and expertise of the contributing authors.

OIMMP Report Authors and Contributors

See Appendix for affiliation abbreviations.

Matthew Anderson	FDEP IRLAP	Kris Kaufman	NOAA
William Arnold	NOAA	Sean King	SWFWMD
Jeff Beal	FWC	Phyllis Klarmann	SFWMD
Anne Birch	TNC	Katie Konchar	FWC
Joshua Breithaupt	UCF	Megan Lamb	ANERR
Aaron Brown	University of Tampa	Jay Leverone	SBEP
Loren D. Coen	FAU	Andrew Lykens	TBW
Allison Conner	CHNEP	Eric Milbrandt	SCCF
Patrice Couch	St. Andrew Bay Watch	Ryan P. Moyer	FWC
Emily Dark	FDEP IRLAP	Melanie Parker	FWC
Katie Davis	FDEP CPAP	William Pine	UF
Nikki Dix	GTMNERR	Kara Radabaugh	FWC
Emma Dontis	FWC	Gary Raulerson	TBEP
Vincent Encomio	UF/IFAS FL Sea Grant	Thomas Ries	ESA
Peter Frederick	UF	Annie Roddenberry	FWC
Stephanie Garvis	UCF	Christi Santi	FWC
Stephen Geiger	FWC	Caitlin Snyder	ANERR
Ray Grizzle	UNH	Mark Thompson	SCCF
Gregory Herbert	USF	Linda Walters	UCF
Erica Hernandez	Alachua Conservation Trust	Amber Whittle	FWC
Stephen Hesterberg	USF	Estelle Wilson	NOAA



Restored oyster reef in the Mosquito Lagoon. Photo credit: Linda Walters.

Executive Summary

Oysters provide a variety of critical ecosystem services to coastal communities in Florida. They improve water quality and clarity as they filter feed, lessen shoreline erosion, and provide a habitat or food source for a wide variety of birds, fish, and invertebrates. Oysters are commercially valuable as a harvested food source, and historically their shell has been mined extensively for construction material. The eastern oyster (*Crassostrea virginica*) is the only reef-building oyster in Florida and forms both subtidal and intertidal reefs. Numerous other species of non-reef-building oysters are less frequent. This report focuses primarily on the eastern oyster, because it is the most abundant oyster in Florida and because it is important as both a keystone species and an ecosystem engineer.

The survival of an oyster reef depends on its shell budget, which is its rate of shell deposition from new oyster growth relative to the rate of shell loss. The rate of growth is limited by basic biological functions of living oysters (rates of growth and reproduction), while the rate of loss is a function of both biotic (e.g., predation, competition) and abiotic (e.g., salinity, temperature, pH) factors that can affect both living oysters and the shells of deceased oysters. Rates of bioerosion, chemical degradation, dis-

solution, and burial affect the length of time the dead shell remains on a reef as viable settlement substrate. The optimum salinity range for eastern oysters is 14 to 28, although they can temporarily tolerate salinity extremes from 5 to 40. Oysters have decreased growth and reproduction at low salinity and can quickly suffer high rates of mortality under freshwater conditions. While oysters can physiologically tolerate high salinity for extended periods of time, in such conditions they are more vulnerable to marine predators, disease, and parasitism. Tolerance of high or low salinity is significantly diminished at high temperatures, which oysters frequently encounter in Florida. Climate change and sea-level rise further alter the frequency and severity of temperature and salinity stress.

Many Florida estuaries have lost 80–90% of the oyster reefs that were present before human development. Altered surface-water flow is one of the major threats to oyster reefs today, as channelization or other mechanisms that concentrate stormwater runoff reduce salinity to levels less than those optimal for oyster survival, growth, and reproduction. Hydrologic alterations, coupled with freshwater withdrawal, also starve downstream areas of freshwater flow, resulting in increased salinity that makes oys-

ters more vulnerable to predation and disease. Significant losses to oyster reefs have occurred due to anthropogenic activities including dredge-and-fill construction, harvesting without substrate replacement, and, in the past, shell mining. Oyster populations must also cope with water pollution, competition with invasive species, sedimentation, and accelerated erosion due to boat wakes. Many of these stressors are increasing as Florida's human population grows. The decline in the Florida oyster population has led to small and often isolated populations spread across Florida's estuaries. Although the long-term and large-scale ramifications of this decline are not well studied, the isolation of these small populations can limit genetic diversity and connectivity between estuaries.

Oyster harvesting is permitted in Florida within designated shellfish harvesting areas. The Florida Department of Agriculture and Consumer Services (FDACS) regulates the opening and closure of these harvesting areas based on health risks to consumers. Both FDACS and the Florida Fish and Wildlife Conservation Commission (FWC) monitor the waters in these shellfish harvesting areas for bacteria, red tide, and chemical pollutants. FWC reports commercial harvesting yields, which have had mandatory reporting since 1986. Historically 90% of the state's harvests originated from Apalachicola Bay in Franklin County; however, harvests from Apalachicola Bay (and consequently statewide harvests) have declined significantly since the 2012–2013 collapse of the bay's fishery.

Large-scale oyster reef mapping relies primarily on georeferenced multispectral or hyperspectral imagery with in situ ground truthing to verify mapping accuracy. Reefs are identified by patterns of color, texture, and shape, but reef identification can be confounded if oysters are intermixed with algae, mud, seagrass, or rubble. Oysters that grow on mangrove roots or seawalls are generally not included in mapping efforts because they are hard to see in aerial imagery. These oysters nevertheless contribute significantly to the oyster population in an estuary. Subtidal reefs can be mapped with side-scan or multibeam sonar or videography with simultaneous acquisition of global positioning system (GPS) data, but ground truthing is necessary to verify the presence of live oysters. Subtidal oyster mapping is complicated by murky water, variable water depth or shallow water, limited benthic relief, and oyster reefs co-occurring with multiple benthic habitats such as seagrass beds and hardbottom. Oyster maps in Florida generally focus on a specific region or estuary. FWC and the Oyster Integrated Mapping and Monitoring Program have compiled available oyster maps across the state, creating the most comprehensive map to date of oyster distribution in Florida (<http://geodata.myfwc.com/datasets/oyster-beds-in-florida>). This map identifies more

than 7,920 ha (19,580 ac) of live oysters, yet gaps remain, and further mapping efforts are needed in several areas.

Oyster monitoring in Florida is conducted by a number of agencies and organizations with a variety of objectives, such as determining the efficacy of hydrologic restoration, the health of oyster fisheries, or the success of restoration efforts, as well as general ecological assessments. While methods used in monitoring programs may vary widely, commonly measured parameters include water quality (salinity, dissolved oxygen, and temperature), reef area, reef height, oyster density, degree of tidal emersion, and oyster size–frequency distribution.

The chapters in this report summarize mapping and monitoring programs for oyster reefs in each region of Florida. Content of each chapter includes a general introduction to the region, location-specific threats to oyster reefs, a summary of selected mapping and monitoring programs, and recommendations for oyster management, mapping, and monitoring. Regional figures include the FWC compilation of oyster maps, FDACS shellfish harvesting areas, and oyster harvesting data from 1951 to 2017.

Through the process of compiling this report and from feedback provided at the OIMMP workshops, several needs and recommendations were identified for Florida oysters:

Management priorities and recommendations

- Manage freshwater flow to mimic natural flow, avoiding rapid salinity changes and prolonged exposure to salinity extremes.
- Add shell and other materials to combat substrate limitation due to extensive harvesting, dredging, or past shell mining. Place substrate on firm sediments to prevent its sinking, and determine ideal locations based on current hydrologic conditions rather than historic reef extent.
- Create and implement a comprehensive oyster fishery management plan that takes into account climate change, variable oyster fishing effort, shell budgets, annually variable freshwater input, and widespread anthropogenic changes in order to prevent overfishing or loss of substrate.
- Replace or supplement hardened shorelines with living shorelines to create habitat and facilitate habitat migration upslope as sea level rises.
- Maintain genetic connectivity of oyster populations between estuaries across the state by rebuilding or maintaining stable oyster populations in all estuaries where they naturally occur.

Mapping priorities and recommendations

- Fill remaining mapping gaps in the Panhandle (Pensacola, Choctawhatchee, and St. Andrew bays), Big Bend and Springs Coast (Apalachee Bay and subtidal oysters), Everglades, and Indian River Lagoon (outside of its major tributaries).
- Complete regular mapping efforts every 5–7 years. Oyster extent is dynamic due to urban development, variable freshwater flow, and changing freshwater management, so maps should be updated regularly.
- Map all oysters, including subtidal oysters and oysters on mangrove roots and seawalls.
- Determine historical extent of oyster reefs to facilitate decision making regarding targets for future reef extent.
- Differentiate between live and dead sections on oyster reefs to track mortality or dead margins on live reefs over time.

Monitoring and research priorities and recommendations

- Conduct standardized and long-term monitoring across multiple estuaries to facilitate comparisons among oyster populations.
- Determine genetic diversity, life history, and habitat characteristics of high-salinity oyster reefs to determine why certain oyster populations survive in high salinity while others are decimated by predators and disease.
- Quantify oyster size structure of oyster populations. Shell height in an oyster population can provide a snapshot of reef resilience because large oysters are disproportionately important to reproductive output and shell budgets.
- Make high-frequency autonomous measurements of temperature and salinity near established oyster reefs in order to capture extreme events such as freshwater pulses, temperature extremes, and hypoxic conditions.