


CHOCTAWHATCHEE BASIN ALLIANCE OF NWF STATE COLLEGE:

Choctawhatchee Bay



Alison McDowell
Director

Choctawhatchee Basin Alliance



Choctawhatchee Bay
is located in the
panhandle of Florida



History

- Florida Dept. of Community Affairs grant used to bring community stakeholders together
- Result of the meetings was the creation of what is now called Choctawhatchee Basin Alliance of NWF State College in 1996
- Quickly became adopted by NWF State College: provided fiduciary stability for grants, contracts, gift-giving, memberships, etc.



Currently—20 years later

- Fully adopted College program
 - space and overhead
 - fiduciary agent
 - support staff (account, purchasing, graphics, etc.)
- Voluntary Community-Based Advisory Committee
- Paid staff
 - Director, 4 Coordinators, 3 PT staff,
- 4 Growing Program areas
 - Monitoring, Restoration, Education, Research





Mission



Choctawhatchee Watershed

- ◆ Watershed area is over 5,000 mi²
- ◆ One main river on east side of watershed
- ◆ Most of the watershed is in Alabama
- ◆ Major land use is Agricultural/ Forest/ Grasslands
- ◆ Most of the urban use is in Okaloosa & Walton Counties



















Water Quality Data







Mean values from 2000-2009







	Western	Middle	Eastern
Number of samples	1074	2586	995
TP ($\mu\text{g/L}$)	12	16	25
TN ($\mu\text{g/L}$)	246	269	413
CHLA ($\mu\text{g/L}$)	4	5	9
Secchi (m)	3	2	1
Color (Pt-Co Units)	9	16	22
Bottom Depth (m)	0.5 - 11.3	0.34 - 11.0	0.30 - 4.88
Surface Salinity (ppt)	22	18	11
Bottom Salinity (ppt)	27	25	17
Surface turbidity (NTU)	4	11	13
Bottom turbidity (NTU)	6	16	18











Summary of Seagrass Abundance

Year	Source	SAV West (Hectares)	SAV Middle (Hectares)	SAV East (Hectares)	Total Calculated (Hectares)	Total Reported (Hectares)
1992	USGS	927	842	2	1,771	1,726
2003	USGS	511	586	2	1,099	1,071
2009	CBA	469	1,008	0	1,477	
2010	CBA	365	1,020	0	1,385	
2011	CBA	190	1,044	0	1,234	



Chain of Eutrophication Models for Assessment of the Potential Nutrient Enrichment of Choctawhatchee Bay, FL, USA

Eutrophication models originally developed for freshwater systems function similarly in Choctawhatchee Bay:

- ◆ Nutrient loading model developed by Canfield and Bachmann (1981) did estimate nutrient concentrations in the bay.
- ◆ There is a significant relation between total phosphorus and chlorophyll in Choctawhatchee bay.
- ◆ Chlorophyll concentration is a significant driver of water clarity in Choctawhatchee Bay
- ◆ Water clarity is not limiting the seagrass coverage in Choctawhatchee Bay

Hoyer, Terrell, Canfield, *Florida Scientist*, 2015



© Scott Jackson - PhotosFlorida.com



Questions?

