

Striped Mud Turtle (Lower Keys Population) Biological Status Review Report

March 31, 2011



**FLORIDA FISH AND WILDLIFE CONSERVATION COMMISSION
620 South Meridian Street
Tallahassee, Florida 32399-1600**

**Biological Status Review
for the
Striped Mud Turtle (Lower Keys Population)
(*Kinosternon baurii*)
March 31, 2011**

EXECUTIVE SUMMARY

The Florida Fish and Wildlife Conservation Commission (FWC) directed staff to evaluate all species listed as Endangered, Threatened or Species of Special Concern as of November 8, 2010 that had not undergone a status review in the past decade. Public information on the status of the Lower Keys population of the striped mud turtle was sought from September 17 through November 1, 2010. A five-member Biological Review Group (BRG) met on November 9-10, 2010. Group members were Bill Turner (FWC lead), Chris Lechowicz (Sanibel-Captiva Conservation Foundation), Peter Meylan (Eckerd College), Paul Moler (independent consultant), and Travis Thomas (University of Florida) (Appendix 1). In accordance with rule 68A-27.0012, Florida Administrative Code (F.A.C.), the BRG was charged with evaluating the biological status of the Lower Keys population of the striped mud turtle using criteria included in definitions in 68A-27.001, F.A.C., and following protocols in the *Guidelines for Application of the IUCN Red List Criteria at Regional Levels (Version 3.0)* and *Guidelines for Using the IUCN Red List Categories and Criteria (Version 8.1)*. Please visit <http://myfwc.com/wildlifehabitats/imperiled/listing-action-petitions/> to view the listing process rule and the criteria found in the definitions.

In late 2010, staff developed the initial draft of this report which included BRG findings and a preliminary listing recommendation from staff. The draft was sent out for peer review and the reviewers' input has been incorporated to create this final report. The draft report, peer reviews, and information received from the public are available as supplemental materials at <http://myfwc.com/wildlifehabitats/imperiled/biological-status/>.

The BRG concluded from the biological assessment that the Lower Keys population of the striped mud turtle met listing criteria. Because the Lower Keys population of the striped mud turtle does not meet the definition of isolated population (significant and discrete population of a species), however, FWC staff recommends that the Lower Keys population of the striped mud turtle not be listed as a Threatened species and that it be removed from the Species of Special Concern list.

This work was supported by a Conserve Wildlife Tag Grant from the Wildlife Foundation of Florida. FWC staff gratefully acknowledges the assistance of the biological review group members and peer reviewers. Staff would like to thank Dale Jackson who compiled the information and wrote the Biological Information section of this report

BIOLOGICAL INFORMATION

Taxonomic Classification –Striped mud turtles from the Lower Keys were formerly considered to be a distinct sub-species (Stejneger 1925, Uzzell and Schwartz 1955). More recent

morphological and molecular studies (Iverson 1978, Lamb and Lovich 1990, Karl and Wilson 2001), however, have indicated that Lower Keys specimens are not sufficiently distinct to justify taxonomic recognition, and most authors do not recognize the Lower Keys population of the striped mud turtle as taxonomically distinct.

Life History and Habitat Requirements – Life history and habitat parameters are summarized range wide by Ernst and Lovich (2009), and for the state of Florida by Wilson et al. (2006). In the Lower Keys, where freshwater habitats are extremely limited, Dunson (1981) captured striped mud turtles in high numbers in small, ephemeral freshwater ponds and brackish water ponds with salinities below 15 ppt. Man-made mosquito control ditches, with longer hydroperiods, also supported high numbers. Rangewide, the species utilizes terrestrial habitats for nesting, migration between ponds (especially males), and aestivation during dry weather (Wilson et al. 2006). In the Lower Keys, the turtles also move onto land to escape drying brackish ponds, which become too saline when water levels recede (Dunson 1992). Though not studied in the Keys, the species' varied diet elsewhere includes insects, worms, snails, algae, seeds, and the remains of vertebrates (e.g., small fishes and amphibians) that are scavenged (Wilson et al. 2006, Ernst and Lovich 2009). Most data about striped mud turtle reproduction have been generated from sites north of the Keys, especially the Florida Peninsula. Iverson (1979) estimated that females reach maturity in about 6 years; males may mature somewhat younger (Wilson et al. 2006). Longevity in the wild is unknown, but based on captive records (Wilson et al. 2006) and data for the closely related *K. subrubrum* (Meshaka and Gibbons 2006), 40 years is a reasonable estimate. Although nesting has been recorded in most months of the year in peninsular Florida, peak activity seems to occur in the fall, with a secondary peak in early summer (Iverson 1979, Wilson et al. 1999, Meshaka and Blind 2001); the extent to which this pattern might be altered in the Lower Keys is unknown. Females may move hundreds of meters from wetlands to nest (Mushinsky and Wilson 1992, Wilson 1996, Wilson et al. 1999); most lay 2-4 clutches of 1-6 eggs each per year (Iverson 1977, Wilson et al. 1999, Meshaka and Blind 2001, Wilson et al. 2006).

Population Status and Trend – Insufficient data are available to document current status and trend quantitatively. The species is known only from mostly small populations (dozens to a few hundred: Dunson 1992) on 11 islands in the Lower Keys (see Geographic Range and Distribution). It can be inferred from the level of development and habitat alteration on these keys that this regional population of the species has declined throughout the 20th century. Perhaps the most resounding example of this is Key West, where Garman (1891) found the species to be “tolerably abundant” in brackish ponds, yet Carr (1940) was unable to find any turtles during the late 1930s and believed the island’s population to have been extirpated. Although construction of mosquito control ditches may have allowed some local populations to increase or recover in terms of numbers (Dunson 1992), this effect could be reversed quickly if some of these ditches are filled.

Geographic Range and Distribution – Striped mud turtles occur throughout Florida including the Florida Keys (Upper Keys, Middle Keys and Lower Keys). The listed Lower Keys population includes only striped mud turtles that occur from the western end of the Seven Mile Bridge to Key West. Specific records of this Lower Keys population, from east to west, are known for Big Pine Key, Little Torch Key, Middle Torch Key, Big Torch Key, Ramrod Key,

Summerland Key, Cudjoe Key, Sugarloaf Key, Johnston Key, Stock Island, and Key West (Florida Natural Areas Inventory 2010).

Quantitative Analyses – Endries et al. (2009) developed a population viability analysis model for the Lower Keys population of the striped mud turtle, but, in the absence of specific microhabitat information, it used overly general habitat criteria that identified 2,539 ha (6,274 ac) of potential habitat. This is a vast (>90%) overestimate, as it includes both pine rockland and tropical hardwood hammock habitats, yet these principally are the upland matrix in which wetlands, the limiting habitat for the species, occur in the Lower Keys. The latter provide no more than ca. 200 ha (500 ac) of potential habitat for the turtle, and this habitat type is significantly threatened by several factors (see Threats) that affect hydroperiod and water quality. The conclusion of the Endries et al. (2009) model, that there is 0% probability of extinction in the next 100 years, is thus untenable and unsupported.

BIOLOGICAL STATUS ASSESSMENT

Threats – The dependence of striped mud turtles on waters of low salinity (< 15 ppt: Dunson 1992) predisposes it to decline and/or extirpation in the Lower Keys. Natural freshwater habitats in the Keys tend to be small (1- 50 acres) and precarious. Regardless of protective measures (regulatory on private lands, natural resource management on public lands), all such water bodies depend upon continued maintenance and protection of natural subsurface freshwater lenses. A myriad of factors associated with development and human habitation threaten these delicate lenses (Lazell 1989), both through direct reduction (hence, recession from the surface) and saltwater intrusion. For mud turtles, creation of artificial mosquito control ditches has partially offset the loss of smaller freshwater bodies, but these ditches do not assure perpetual habitat. Perhaps the most serious threat to all freshwater and brackish habitats in the Keys is sea level rise that is predicted to occur as a direct consequence of global warming (Field et. al. 2007). Because the striped mud turtle inhabits only a few islands in the Lower Keys (very small Extent of Occurrence and Area of Occupancy), the Lower Keys population is naturally vulnerable to threats by stochastic events. Although the species has survived many hurricanes, severe saltwater over wash from very large storms has the potential to increase salt content of fresh and brackish water ponds to an extent that would eliminate them as suitable habitat for the mud turtle (Dunson 1992). Random events of severe pollution also provide a serious threat, as exemplified by the 2010 (MC 252) oil spill in the Gulf of Mexico. Protective booms or other measures would probably not prevent oiled waters from being cast over the entire Keys during a large hurricane; such a disaster would likely extirpate many local and regional populations of freshwater life, including mud turtles and their prey. Although not studied, it is likely that predators, particularly raccoons, take high percentages of mud turtle eggs as well as surviving young as is the case with most turtles. This reduces the potential for already small, isolated populations of these turtles to recover from declines caused by any factors.

Lower Keys Population Assessment – Findings from the BRG are included in the Biological Status Review Information Findings tables, below.

LISTING RECOMMENDATION

The BRG concluded from the biological assessment that the Lower Keys population of the striped mud turtle met listing criteria. Although the Lower Keys population of the striped mud turtle may be somewhat isolated from striped mud turtles to the north by an expanse of salt water, staff did not believe the population was sufficiently distinct to warrant listing as an isolated population (significant and discrete population of a species). Therefore, FWC staff recommends the Lower Keys population of the mud turtle not be listed as a Threatened species and that it be removed from the Species of Special Concern list.

SUMMARY OF THE INDEPENDENT REVIEW

Dr. J. Whitfield Gibbons and Dr. John B. Iverson provided peer review of this report. Both reviewers agreed that the review was thorough and supported the findings of the BRG. Although both were concerned about striped mud turtle population in the Lower Keys, they concurred that it did not meet the criteria for an isolated population and supported staff's recommendation to remove the population from the list of Threatened species. Dr. Iverson provided additional information about generation time and pointed out that the warmer climate in the Keys may shorten it. Using a shorter generation time, however, would not have changed the findings of the assessment.

LITERATURE CITED

- Carr, A. F. 1940. A contribution to the herpetology of Florida. University of Florida Publications, Biological Sciences Series 3:1-118.
- Dunson, W. A. 1981. Behavioral osmoregulation in the key mud turtle, *Kinosternon b. baurii*. Journal of Herpetology 15:163-173.
- Dunson, W. A. 1992. Striped mud turtle. Pages 105-110 in P. E. Moler, editor. Rare and endangered biota of Florida, vol. III: amphibians and reptiles. University Press of Florida, Gainesville, Florida, USA. 291pp.
- Endries, M., B. Stys, G. Mohr, G. Kratimenos, S. Langley, K. Root, and R. Kautz. 2009. Wildlife habitat conservation needs in Florida. Fish and Wildlife Research Institute Technical Report TR-15. 178 pp.
- Ernst, C. H., and J. E. Lovich. 2009. Turtles of the United States and Canada. Second edition. The Johns Hopkins University Press, Baltimore, Maryland, USA. 827pp.
- Field, C.B., L.D. Mortsch, M. Brklacich, D.L. Forbes, P. Kovacs, J.A. Patz, S.W. Running and M.J. Scott, 2007: North America. *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds., Cambridge University Press, Cambridge, UK, 617-652.
- Florida Natural Areas Inventory. 2010. Tallahassee, Florida, USA. [contacts: via the Internet at www.fnai.org, or telephone 850-224-8207]
- Garman, S. 1891. On a tortoise found in Florida and Cuba, *Cinosternum baurii*. Bulletin of the Essex Institute 23:141-144.
- Iverson, J. B. 1977. Reproduction in freshwater and terrestrial turtles of north Florida. Herpetologica 33:205-212.
- Iverson, J. B. 1978. Variation in striped mud turtles, *Kinosternon baurii* (Reptilia, Testudines, Kinosternidae). Journal of Herpetology 12:135-142.
- Iverson, J. B. 1979. The female reproductive cycle in north Florida *Kinosternon baurii* (Testudines: Kinosternidae). Brimleyana 1:37-46.
- Karl, S. A., and D. S. Wilson. 2001. Phylogeography and systematics of the mud turtle, *Kinosternon baurii*. Copeia 2001:797-801.
- Lamb, T., and J. E. Lovich. 1990. Morphometric validation of the striped mud turtle (*Kinosternon baurii*) in the Carolinas and Virginia. Copeia 1990:613-618.

- Lazell, J. D., Jr. 1989. Wildlife of the Florida Keys: a natural history. Island Press. Washington, D.C. 254pp.
- Meshaka, W. E., Jr., and E. Blind. 2001. Seasonal movements and reproduction in the striped mud turtle (*Kinosternon baurii*) from the southern everglades, Florida. *Chelonian Conservation and Biology* 4:75-80.
- Meshaka, W. E., and J. W. Gibbons. 2006. *Kinosternon subrubrum* – eastern mud turtle. Pages 189-196 in P. A. Meylan, editor. Biology and conservation of Florida turtles. *Chelonian Research Monographs* No. 3.
- Mushinsky, H. R., and D. S. Wilson. 1992. Seasonal occurrence of *Kinosternon baurii* on a sandhill in central Florida. *Journal of Herpetology* 26:207-209.
- Stejneger, L. 1925. New species and subspecies of American turtles. *Journal of the Washington Academy Sciences* 15:462-463.
- Uzzell, T. M., Jr., and A. Schwartz. 1955. The status of the turtle *Kinosternon bauri palmarum* Stejneger with notes on variation in the species. *Journal of the Elisha Mitchell Society* 71:28-35.
- Wilson, D. S. 1996. Nesting ecology and nest site selection in the striped mud turtle, *Kinosternon baurii*, in central Florida. Ph.D. dissertation, University of South Florida, Tampa, Florida.
- Wilson, D. S., H. R. Mushinsky, and E. D. McCoy. 1999. Nesting behavior of the striped mud turtle, *Kinosternon baurii* (Testudines: Kinosternidae). *Copeia* 1999:958-968.
- Wilson, D. S., H. M. Mushinsky, and E. D. McCoy. 2006. *Kinosternon baurii* – striped mud turtle. Pages 180-188 in P. A. Meylan, editor. Biology and conservation of Florida turtles. *Chelonian Research Monographs* No. 3.

Biological Status Review Information
Findings

Species/taxon: Lower Keys population of the striped mud turtle

Date: November 9-10, 2010

Assessors: Chris Lechowicz, Peter Meylan, Paul Moler,
Bill Turner and Travis Thomas

Generation length: 17.5

Criterion/Listing Measure	Data/Information	Data Type*	Sub-Criterion Met?	References
*Data Types - observed (O), estimated (E), inferred (I), suspected (S), or projected (P). Sub-Criterion met - yes (Y) or no (N).				
(A) Population Size Reduction, ANY of				
(a)1. An observed, estimated, inferred or suspected population size reduction of at least 50% over the last 10 years or 3 generations, whichever is longer, where the causes of the reduction are clearly reversible and understood and ceased ¹	insufficient data	S	N	
(a)2. An observed, estimated, inferred or suspected population size reduction of at least 30% over the last 10 years or 3 generations, whichever is longer, where the reduction or its causes may not have ceased or may not be understood or may not be reversible ¹	Inferred from habitat loss. Lower Keys have experienced extensive development with reduction of natural freshwater habitats in last half century. Populations on Key West and Stock Island likely highly reduced or eliminated.	I	Y	Dunson 1992
(a)3. A population size reduction of at least 30% projected or suspected to be met within the next 10 years or 3 generations, whichever is longer (up to a maximum of 100 years) ¹	This is possible but there is little information to go on; perhaps can obtain growth projections from Monroe County. Much remaining habitat protected in National Key Deer Refuge.	I	N	
(a)4. An observed, estimated, inferred, projected or suspected population size reduction of at least 30% over any 10 year or 3 generation period, whichever is longer (up to a maximum of 100 years in the future), where the time period must include both the past and the future, and where the reduction or its causes may not have ceased or may not be understood or may not be reversible. ¹	Highly likely, given patterns of past & projected development, alteration of natural habitats, sea level rise, salt water intrusion, decline in freshwater lens, hurricanes, other stressors	S	Y	Dunson 1992
¹ based on (and specifying) any of the following: (a) direct observation; (b) an index of abundance appropriate to the taxon; (c) a decline in area of occupancy, extent of occurrence and/or quality of habitat; (d) actual or potential levels of exploitation; (e) the effects of introduced taxa, hybridization, pathogens, pollutants, competitors or parasites.				
(B) Geographic Range, EITHER				
(b)1. Extent of occurrence < 20,000 km ² (7,722 mi ²) OR	348 km ² , estimated area of 11 keys	E	Y	D. Jackson generated GIS polygon from Florida Natural Areas Inventory (FNAI) records
(b)2. Area of occupancy < 2,000 km ² (772 mi ²)	<20 km ² , limited freshwater habitats, many artificial ditches, ponds.	E	Y	GIS habitat analysis by FWC/Stys

AND at least 2 of the following:				
a. Severely fragmented or exist in ≤ 10 locations	Fragmented-11 keys (small islands), naturally severely fragmented by intervening salt water (ocean), rare accidental transport possible but not significant.	O	Y	FNAI data, Dunson 1992
b. Continuing decline, observed, inferred or projected in any of the following: (i) extent of occurrence; (ii) area of occupancy; (iii) area, extent, and/or quality of habitat; (iv) number of locations or subpopulations; (v) number of mature individuals	Projected decline, all categories, with loss of natural freshwater habitat related to development, sea level rise, stochastic events, fire ants	S	Y	Dunson 1992, Forys (Marsh rabbit info.)
c. Extreme fluctuations in any of the following: (i) extent of occurrence; (ii) area of occupancy; (iii) number of locations or subpopulations; (iv) number of mature individuals	No; extreme fluctuations unlikely in long-lived species;	O	N	
(C) Population Size and Trend				
Population size estimate to number fewer than 10,000 mature individuals AND EITHER	Likely less than 10,000 Even the densest known population (Summerland Key) is estimated to be in the hundreds; estimated 50 on undisturbed Johnson Key	S	Y	Dunson 1992
(c)1. An estimated continuing decline of at least 10% in 10 years or 3 generations, whichever is longer (up to a maximum of 100 years in the future) OR	see above Criterion A	S	Y	
(c)2. A continuing decline, observed, projected, or inferred in numbers of mature individuals AND at least one of the following:		S	Y	
a. Population structure in the form of EITHER				
(i) No subpopulation estimated to contain more than 1000 mature individuals; OR	Even the densest known population (Summerland Key) is estimated to be in the hundreds.	E	Y	Dunson 1992
(ii) All mature individuals are in one subpopulation	known from >10 islands	O	N	Dunson 1992, FNAI
b. Extreme fluctuations in number of mature individuals	No; extreme fluctuations unlikely in long-lived species	O	N	
(D) Population Very Small or Restricted, EITHER				
(d)1. Population estimated to number fewer than 1,000 mature individuals; OR	Uncertain, probably >1000-2000	E	N	Dunson 1992, FNAI
(d)2. Population with a very restricted area of occupancy (typically less than 20 km ² [8 mi ²]) or number of locations (typically 5 or fewer) such that it is prone to the effects of human activities or stochastic events within a short time period in an uncertain future	<20 km ² , very restricted, remnant wetlands and drainage ditches (insert estimate if available)	S	Y	GIS habitat analysis by FWC/Stys
(E) Quantitative Analyses				

e1. Showing the probability of extinction in the wild is at least 10% within 100 years	No appropriate models	S	N	
Initial Finding (Meets at least one of the criteria/sub-criteria OR Does not meet any of the criteria/sub-criteria)	Reason (which criteria/sub-criteria are met)			
Meets multiple criteria	A2+A4; B1+B2ab (i,ii,iii,iv,v); C1+C2a(i); D2			
Is species/taxon endemic to Florida? (Y/N)	N			
If Yes, your initial finding is your final finding. Copy the initial finding and reason to the final finding space below. If No, complete the regional assessment sheet and copy the final finding from that sheet to the space below.				
Final Finding (Meets at least one of the criteria/sub-criteria OR Does not meet any of the criteria/sub-criteria)	Reason (which criteria/sub-criteria are met)			
Meets multiple criteria	A2+A4; B1+B2ab (i,ii,iii,iv,v); C1+C2a(i); D2			

Regional Assessment

1	<p align="center">Biological Status Review Information</p> <p align="center">Regional Assessment</p>	Species/taxon:	Lower Keys population of the striped mud turtle
2		Date:	November 9- 10, 2010
3		Assessors:	Chris Lechowicz, Peter Meylan, Paul Moler,
4			Bill Turner and Travis Thomas
5			
6			
7			
8	Initial finding		
9			
10	2a. Is the species/taxon a non-breeding visitor? (Y/N/DK). If 2a is YES, go to line 18. If 2a is NO or DO NOT KNOW, go to line 11.		N
11	2b. Does the Florida population experience any significant immigration of propagules capable of reproducing in Florida? (Y/N/DK). If 2b is YES, go to line 12. If 2b is NO or DO NOT KNOW, go to line 17.		N
12	2c. Is the immigration expected to decrease? (Y/N/DK). If 2c is YES or DO NOT KNOW, go to line 13. If 2c is NO go to line 16.		
13	2d. Is the regional population a sink? (Y/N/DK). If 2d is YES, go to line 14. If 2d is NO or DO NOT KNOW, go to line 15.		
14	If 2d is YES - Upgrade from initial finding (more imperiled)		
15	If 2d is NO or DO NOT KNOW - No change from initial finding		
16	If 2c is NO or DO NOT KNOW - Downgrade from initial finding (less imperiled)		
17	If 2b is NO or DO NOT KNOW - No change from initial finding		No change
18	2e. Are the conditions outside Florida deteriorating? (Y/N/DK). If 2e is YES or DO NOT KNOW, go to line 24. If 2e is NO go to line 19.		
19	2f. Are the conditions within Florida deteriorating? (Y/N/DK). If 2f is YES or DO NOT KNOW, go to line 23. If 2f is NO, go to line 20.		
20	2g. Can the breeding population rescue the Florida population should it decline? (Y/N/DK). If 2g is YES, go to line 21. If 2g is NO or DO NOT KNOW, go to line 22.		
21	If 2g is YES - Downgrade from initial finding (less imperiled)		
22	If 2g is NO or DO NOT KNOW - No change from initial finding		
23	If 2f is YES or DO NOT KNOW - No change from initial finding		
24	If 2e is YES or DO NOT KNOW - No change from initial finding		
25			
26	Final finding		No change

Additional notes -- Calculation of generation time presented at the BSR group meeting

Generation length is defined as the average age of parents of the current cohort, which is greater than the age at first breeding and less than the age of the oldest breeding individual. We estimate generation length for the Lower Keys population of the striped mud turtle as follows. Age to maturity is estimated at a mean of 5 years based on Iverson (1979) and Wilson et al. (2006). Longevity is estimated at ca. 40 years maximum based on data from Wilson et al. (2006) and the closely related *K. subrubrum* (Meshaka and Gibbons 2006). There is no reason to distinguish sexes. 30 years may be a reasonable life expectancy for most mature individuals. Generation length is estimated as $(5 + 30)/2 = 17.5$ years.

APPENDIX 1. Brief biographies of the Striped mud turtle Biological Review Group members.

Chris Lechowicz is the Interim Director of the Wildlife Habitat Management Program and staff herpetologist at the Sanibel-Captiva Conservation Foundation where he has worked since 2002. He has a B.S. in Zoology and Computer Science from Southern Illinois University at Carbondale and will complete his M.S. in Environmental Science from Florida Gulf Coast University in 2010. Chris's focus is on riverine turtles with a specialty on the Genus *Graptemys*. Chris is a member of the IUCN/SCC Tortoise and Freshwater Turtle Specialists Group as well as a board member of the Florida Turtle Conservation Trust.

Dr. Peter A. Meylan received his Ph.D. from the University of Florida. He is a Professor of Biology at Eckerd College in Saint Petersburg, FL. His research interests include the evolutionary history, ecology, and conservation biology of amphibians and reptiles, especially turtles. Current research includes 2 sea turtle projects: an investigation of the ecology and migrations of sea turtles of Bocas del Toro Province, Panama (funded by the Wildlife Conservation Society) and the Bermuda Turtle Project, which is a cooperative project with the Bermuda Aquarium and the Caribbean Conservation Corporation (as well as continuing to work with Florida freshwater turtles with the Eckerd Herpetology Club on the Rainbow River). He has many scientific articles on turtles and is the editor of a book on the biology and conservation of Florida turtles.

Paul E. Moler received his M.S. in Zoology from the University of Florida in 1970 and his B.A. in Biology from Emory University in 1967. He retired in 2006 after working for 29 years as a herpetologist with FWC, including serving as administrator of the Reptile and Amphibian Subsection of the Wildlife Research Section. He has conducted research on the systematics, ecology, reproduction, genetics, and conservation biology of a variety of herpetofaunal species in Florida, with primary emphasis on the biology and management of endangered and threatened species. He served as Chair for the Florida Committee on Rare and Endangered Plants and Animals in 1992–94, Chair of the Committee on Amphibians and Reptiles since 1986, and editor of the 1992 volume on amphibians and reptiles. Paul has more than 90 publications on amphibians and reptiles.

Travis Thomas is a graduate student in Wildlife Ecology and Conservation at the University of Florida. His research primarily focuses on the ecology and management of macrofauna in riparian systems. He received his Bachelor's Degree in Natural Resources Conservation from the University of Florida in May 2009. He has worked for 7 years on *M. temminckii* and most recently worked on gopher tortoises for FWC under Joan Berish. He worked for 3 years in the Herpetology Dept. under Dr. Kenneth Krysko at the Florida Museum of Natural History. He has spent time as a volunteer on numerous projects in Kenya, Africa, under the supervision of Leigh Ecclestone and the Kenyan Wildlife Service. He has published several notes on the ecology and distribution of reptiles and is currently a co-author on a study of the ecology of *M. temminckii* in O'Leno State Park as well as the primary author on a study of the morphology of *M. temminckii*.

William M. Turner received his B.S. from Erskine College and M.S. in Biology from the University of South Alabama. From 2003 to 2007, he was the Herpetological Coordinator for the Wyoming Game and Fish Department. In Wyoming, he conducted statewide surveys for

amphibians and reptiles, focusing on emerging amphibian diseases and the impacts of resource development on native reptiles. Since 2007, he has been the Herp. Taxa Coordinator for FWC in the Division of Habitat and Species Conservation. He has conducted research on native amphibians and reptiles in Florida, Alabama and Wyoming that resulted in several published papers and reports.

APPENDIX 2. Summary of letters and emails received during the solicitation of information from the public period of September 17, 2010 through November 1, 2010.

No additional public information was received during the public solicitation period.